

## Carbon and its Compounds

### Introduction


---

- Compounds of carbon are known as organic compounds.
- All organic compounds contain hydrogen along with carbon.
- Since, they are the fundamental organic compounds, they are also known as hydrocarbons.
- The study of carbon compounds such as hydrocarbons and its derivatives is called organic chemistry.


### Bonding in Carbon

---

- Carbon atom has four electrons in its outermost shell.



- It requires four electrons to achieve the stable, 8 electron, inert gas configuration.
- Carbon atoms can achieve the inert gas electron arrangement only by sharing their electrons. Hence, carbon always forms covalent bonds.
- The valency of carbon is four since one carbon requires 4 electrons to achieve the nearest inert gas configuration. Thus, we can say that carbon is tetravalent.
- The four valencies of carbon are usually represented by drawing four short lines around the symbol of carbon (C).




### Allotropes of Carbon

- The various physical forms in which an element can exist are called the allotropes of that element.
- Carbon has three allotropes:
  - Diamond
  - Graphite
  - Buckminster fullerene


#### Diamond

- In diamond, each carbon atom is bonded to four other carbon atoms, forming a three dimensional structure.
- The rigid structure of diamond makes it a very hard substance.
- It is a non-conductor of electricity since there are no free electrons in a diamond crystal.
- It can be synthesised by subjecting pure carbon to a very high pressure and temperature.



### Graphite

- In graphite, each carbon atom is bonded to three other carbon atoms in the same plane, giving a hexagonal array.
- One of the bonds is a double bond and thus the valency of carbon is satisfied.
- Graphite structure is formed by the hexagonal arrays being placed in layers, one above another.
- Graphite is smooth and slippery.
- It is a very good conductor of electricity due to the presence of free electrons.



### Fullerene

- It is an allotrope of carbon containing clusters of 60 carbon atoms joined together to form spherical molecules.
- There are 60 carbon atoms in a molecule of buckminsterfullerene, so its formula is  $C_{60}$ .
- The allotrope was named buckminsterfullerene after the American architect Buckminster Fuller.




## Versatile Nature of Carbon

The two characteristic properties of the element carbon which leads to the formation of a very large number of organic compounds are:


- i. **Catenation:** The property of the element carbon due to which its atoms can join one another to form long carbon chains is called catenation.

### Types of Chains


- Straight chain
  - Branched chains
  - Closed or ring chains
- a. Straight chain of carbon atoms



- b. Branched chain of carbon atoms



- c. Closed or ring chain of carbon atoms




- ii. **Tetravalency:** Carbon has a valency of four. So, it is capable of bonding with four other atoms of carbon or atoms of some other monovalent element.

Compounds of carbon are formed with oxygen, nitrogen, hydrogen, sulphur, chlorine and many other elements, giving rise to compounds with specific properties which depend on the elements other than the carbon present in the molecule.

## Classification of Hydrocarbons

### Saturated Hydrocarbons

- Hydrocarbons in which the carbon atoms are connected by only single bonds are called saturated hydrocarbons.
- Saturated hydrocarbons are called alkanes.
- General formula of alkanes:  $C_nH_{2n+2}$ ,  $n$  = number of carbon atoms
- Methane and ethane are saturated hydrocarbons, which contain only carbon-carbon single bonds.




### Unsaturated Hydrocarbons (Alkenes and Alkynes)

- Hydrocarbons in which two carbon atoms are connected by a double or a triple bond are called unsaturated hydrocarbons.
- Unsaturated hydrocarbons are of two types
  - Alkenes
  - Alkynes

- Alkenes**

An unsaturated hydrocarbon in which two carbon atoms are connected by a double bond is called an alkene.

Alkenes contain the  group.

General formula:  $C_nH_{2n}$ , where  $n$  = number of carbon atoms

- Alkynes**


An unsaturated hydrocarbon in which two carbon atoms are connected by a triple bond is called an alkyne.

An alkyne contains the  $-C \equiv C-$  group.

General formula:  $C_nH_{2n-2}$ , where  $n$  = number of carbon atoms


## Cyclic Hydrocarbons

- Hydrocarbons in which the carbon atoms are arranged in the form of a ring are called cyclic hydrocarbons.
- Cyclic hydrocarbons may be saturated or unsaturated.
  - Saturated cyclic hydrocarbon**
    - Cyclohexane is an example of a saturated cyclic hydrocarbon.
    - Formula:  $C_6H_{12}$
    - Cyclohexane contains 6 carbon atoms arranged in a hexagonal ring, with each carbon atom attached to 2 hydrogen atoms.



- Unsaturated cyclic hydrocarbon**

- Benzene is an example of an unsaturated cyclic hydrocarbon.
- Formula:  $C_6H_6$
- Benzene is made up of 6 carbon atoms and 6 hydrogen atoms.



## Functional Groups

- All organic compounds are derivatives of hydrocarbons.
- Derivatives are obtained by replacing one or more hydrogen atoms by some other atom or group of atoms.
- The new set of compounds formed after replacement has functions different from the parent hydrocarbon.
- Functional group:** An atom or a group of atoms present in the molecules, which determines the characteristics property of the organic compounds, is called the functional group.

Functional group	General formulae	Organic compound	Suffix	Examples with common & IUPAC name
Halide-X (F,Cl,Br,I)	R-X	Haloalkanes	-ane	CH <sub>3</sub> Cl Common name: Methyl chloride IUPAC name: Chloromethane
Hydroxyl-OH	R-OH	Alcohols	-ol	C <sub>2</sub> H <sub>5</sub> OH Common name : Ethyl alcohol IUPAC name: Ethanol
Aldehyde- CHO	$\begin{array}{l} \text{H} \\ \diagdown \\ \text{C}=\text{O} \\ \diagup \\ \text{R} \end{array}$	Aldehydes	-al	CH <sub>3</sub> CHO Common name: Acetaldehyde IUPAC name: Ethanal
Carboxyl- COOH	$\begin{array}{c} \text{O} \\ \parallel \\ \text{R}-\text{C}-\text{O}-\text{H} \end{array}$	Carboxylic acids	-oic acid	CH <sub>3</sub> CH <sub>2</sub> COOH Common name: Propionic acid IUPAC name: Propanoic acid
Keto $\begin{array}{c} \text{O} \\ \parallel \\ -\text{C}- \end{array}$	$\begin{array}{c} \text{O} \\ \parallel \\ \text{R}-\text{C}-\text{R}' \end{array}$	Ketones	-one	CH <sub>3</sub> COC <sub>2</sub> H <sub>5</sub> Common name: Diethyl ketone IUPAC name: Pentanone
Ethers $\begin{array}{c}   \quad   \\ -\text{C}-\text{O}-\text{C}- \\   \quad   \end{array}$	R-O-R'	Ethers	-oxy	CH <sub>3</sub> -O-C <sub>2</sub> H <sub>5</sub> Common name: Ethyl methyl ether IUPAC name: Methoxy ethane

## Homologous Series

---



It is a group of organic compounds having a similar structure and chemical properties in which the successive compounds differ by a  $-\text{CH}_2$  group.

### Characteristics of a Homologous Series

- Each member of the series differs from the preceding one by the addition of a  $-\text{CH}_2$  group and by 14 a.m.u.
- All members of a homologous series have the same general formula.
- The physical properties of the members show a gradation in properties as their molecular mass increases.
- The chemical properties also show a gradient similarity.

**For example:** the general formula for alkane is  $\text{C}_n\text{H}_{2n+2}$  and alkene is  $\text{C}_n\text{H}_{2n}$ .

**For example:** Methane and ethane react with chlorine to form methyl chloride and ethyl chloride respectively.



- All members of a homologous series can be prepared by the same general method of preparation.

**For example:** Alcohols can be prepared from alkyl halides.

